



UV and X-ray Emission from Impacts of Fragmented Accretion Streams on Classical T Tauri stars

S. Colombo¹, S. Orlando¹, G. Peres^{1, 2}, C. Argiroffi^{1, 2}, F. Reale^{1, 2}

¹ INAF - Osservatorio Astronomico di Palermo, Piazza del Parlamento 1, 90143 Palermo, Italy
² Dipartimento di Fisica e Chimica, Università degli Studi di Palermo, Piazza del Parlamento 1, 90143, Palermo, Italy



Context

The accretion process in Classical T Tauri Stars (CTTSs) can be studied through the analysis of some UV and X-ray emission lines which trace hot gas (Sacco et al. 2008, Orlando et al. 2010).

In the UV band these lines consist of multiple components of plasma with different Doppler shifts whose origin is not clear (Ardila et al. 2013).

In this work we investigate the origin of UV and X-ray emission arising from regions where accretion streams impact onto the stellar chromosphere (Reale et al. 2013) and if and how the stream fragmentation determines the observed asymmetries and redshift of UV emission lines in CTTSs.

Model

We model the impact of a fragmented accretion stream onto the chromosphere of a CTTS with 2D axisymmetric magneto-hydrodynamic simulations. Our model includes:

- Stellar magnetic field
- Gravity
- Thermal conduction
- Radiative cooling from an optically thin plasma

From the model results, we synthesize the UV and X-ray emission and its Doppler shift along the line of sight.

The simulations were performed using the PLUTO code (Mignone et al. 2007).




Figure 2: Evolution of density (left half-panel) and temperature (right half-panel) in log scale of plasma for randomly fragmented stream. White lines represent magnetic field lines (Colombo et al. 2016). Note the complex structure of impact region due to stream random fragmentation.

Results and Conclusion

The impacts of accreting blobs onto stellar chromosphere produce reverse shock propagating through the blobs and shocked upflows. These upflows hit and shock back the next downfalling fragments (see Fig. 1 and 2).

As a result:

- Different velocities, densities and temperatures coexist in the shock region
- In each component of the CIV (1550 Å) doublet we see a slow component (SC) shifted at 50 km s^{-1} and a fast component (FC) with redshift ranging between 200 and 400 km s^{-1} . The former is due to post-shock plasma which cools down and decelerates at the base of the accretion column, the latter originates from plasma which cools down due to thermal instabilities and from the interaction of upflowing surges with downfalling blobs (See Fig. 3)
- Our simulations show that the relative intensity of the FC increases with the stream fragmentation
- Each line of OVIII (18.97 Å) doublet is redshifted to speed of 150 km s^{-1}

Our model predicts CIV lines profile remarkably similar to the observed ones and explains their origin in a natural way as due to stream fragmentation.

References

- Ardila et al. (2013), Hot Gas Lines in T Tauri Stars, *ApJS*, 207:1.
- Colombo et al. (2016), Impacts of fragmented accretion stream onto classical T Tauri stars: UV and X-ray emission lines, *A&A* (submitted)
- Mignone et al. (2007), PLUTO: A Numerical Code for Computational Astrophysics, *ApJS*, 170:228-242.
- Orlando et al. (2010), X-ray emitting MHD accretion shock in classical T Tauri stars. Case for moderate to high plasma- β values, *A&A*, 510:A71.
- Reale et al. (2013), Bright Hot Impacts by Erupted Fragments Falling Back on the Sun: A Template for Stellar Accretion, *Science*, 341:251-253.
- Sacco et al. (2008), X-ray emission from dense plasma in classical T Tauri stars: hydrodynamic modeling of the accretion shock. *A&A*, 491:L17-L20.

Acknowledgment: We acknowledge support from INAF through the Progetto Premiale: "A Way to Other Worlds" of the Italian Ministry of Education, University, and Research. The CINECA Award HP10CW941 and the HPC facility (SCAN) of the INAF – Osservatorio Astronomico di Palermo, for the availability of high performance computing resources and support. PLUTO is developed at the Turin Astronomical Observatory in collaboration with the Department of Physics of Turin University. CHIANTI is a collaborative project involving the NRL (USA), the Universities of Florence (Italy) and Cambridge (UK), and George Mason University (USA).




Figure 1: Evolution of density (left half-panel) and temperature (right half-panel) in log scale of plasma for the reference case of a train of blob. White lines represent magnetic field lines (Colombo et al. 2016).




Figure 3: Synthetic profiles of CIV (left) and OVIII doublets (right); The top panel compares our reference simulation of a train of blobs (TRAIN-REF) with the case of a continuous stream (CONT) and with the case of a train of blobs with higher fragmentation (TRAIN-FRAG). The bottom panels show profiles from simulations of a randomly fragmented stream. The dotted grey lines mark the rest positions of the lines of the doublet. All the profiles are normalized to their maximum value. (Colombo et al. 2016)